

“TURPRO”

Improving productivity on turbot farms

Land-based aquaculture is an increasingly important sector within the fisheries industry. There has been a move towards farming a greater diversity of marine fish species, including cod, halibut and sea bass. New species promote growth in the aquaculture sector, and address an increasing consumer demand for farmed fish.

Turbot is a large marine flatfish that has great potential in land-based aquaculture. It is native to Europe, difficult to catch at sea, is esteemed for its flesh, and has a high market value. However, technological improvements are needed to exploit turbot's potential.

The co-operative research project **TURPRO** aimed to **enhance productivity throughout the turbot production chain, from juvenile fish to the market**. The project studied the impact of temperature, photo-period and water quality – key rearing factors – on growth performance, food conversion efficiency, age at first maturity and fish welfare, so as to enable a year-round supply of juveniles reared in intensive land-based systems at high densities. In addition, the impact of processing methods (e.g. slaughtering and storage), especially designed for turbot, on flesh quality and fish yield was investigated.

Temperature

Temperature is an important factor determining turbot growth rates. The effect of step wise temperature regulation was studied as compared to a constant temperature regime, whereby juvenile turbot were reared at two different temperatures (16° and 20°C). After two months, part of the fish were switched (some went from 20° to 16° and others from 16° to 20°) and the other part continued to be reared at the same temperature, which yielded four sub-groups. The 20°-16° group gained 5-11% more weight over

the same period compared to the other three groups.

Photo-period

Extending photo-period is a means of delaying sexual maturity, which, in turn, is desirable to maximise growth. Juvenile turbot were reared at four different photoperiods, and although prolonged exposure to continuous light had a positive effect on the incidence of maturation, it did not have the same impact on growth. More experiments would therefore be necessary to establish the optimum turbot photo-period.

Water quality

The benefits of environmental manipulation of fish growth are reduced in systems with poor water quality. Maintaining water quality is essential in recirculation stocking densities with high stocking densities. Laboratory experiments with levels of ammonia and dissolved oxygen showed that chronic low and high Total Ammonia Nitrogen (TAN) reduced growth, but mild hyperoxia (oxygen overload) appeared to counteract the negative effects of this chronic exposure. Abrupt and limited peaks of TAN reduced growth significantly. Water renewal rates (WRR) were also tested and growth appeared to improve with

Project acronym:

TURPRO

Full title of Project:

Biological optimisation and development of processing methods for turbot farming

EU contract number:

508070

Web-site:

[pi.ijs.si/pibrain.exe?
Cm=Project&Project=TURPRO&Reference=508070](http://pi.ijs.si/pibrain.exe?Cm=Project&Project=TURPRO&Reference=508070)

Coordinator

Mr. Albert Imsland
Akvaplan-NIVA AS
Polar Environment Centre
9296 Tromsø
Norway

Phone: +354 562 58 00

Fax: +354 552 88 01

Email:

albert.imsland@akvaplan.niva.no

“LOWER COSTS AND
HIGHER YIELDS WILL
ENABLE SMES TO EXPAND
PRODUCTION AND ENSURE
THAT EUROPE REMAINS A
WORLD LEADER IN TURBOT
FARMING”

WRR, but not significantly. Underlying mechanisms improving growth could not be explained by different levels of the water quality parameters analysed; it was therefore concluded that they are probably the sum of all water quality parameters.

Flesh quality

The flesh quality in turbot appeared to be highly affected by the season: fish slaughtered in November had a textural shear force which was approximately twice as hard as fast growing fish slaughtered in June. Quality was also affected by the stunning method (percussive stunning, electrical stimulation and exsanguination on ice) and following stress. When studying the processing methods, like gill bleeding, gutting, pre rigor filleting and freezing, it was found that the colour of the fillets can be tuned by the (post) slaughter conditions and that non-bleeding conditions (no gutting, no exsanguinations) result in a more crème colour of the fillet, which is more like the desired fillet colour of wild turbot.

Market opportunities

In the final phase of the project, the market opportunities for farmed turbot were determined and compared to those for wild turbot. In general the wild turbot appeared to be less firm and more tender but also less sticky

compared to farmed turbot. As to freshness, it was found that the shelf-life of farmed turbot can be seven days longer than the wild species. A consumer study showed that consumers attached most value to freshness, price, and quality or labelling.

Turbot has a very high market potential. Although popular as a table fish in France and Spain, it is only just being discovered in northern Europe. It attracts high prices and fish farmers can make good profits from it. Production costs are nevertheless high for smaller European enterprises, and competition from China is increasing fast. **The protocol produced by TURPRO, in the form of a practical manual, will help to reduce costs and increase productivity on turbot farms in a sustainable and environmentally friendly manner.**